GAD-7 Screening Questions^{1,2}

During the last 2 weeks, how often have you been bothered by the following problems?

Problem	Not at all	Several days	More than half the days	Nearly every day	
1. Feeling nervous, anxious, or on edge	0	1	2	3	
2. Not being able to stop or control worrying	0	1	2	3	
3. Worrying too much about different things	0	1	2	3	
4. Trouble relaxing	0	. 1	2	3	
5. Being so restless that it is hard to sit still	0	1	2	3	
6. Becoming easily annoyed or irritable	0	1	2	3	
7. Feeling afraid as if something awful might happen	0	1	2	3	
	Total each column:				
		Total score (Add columns 2,3,4):			

Adapted from Spitzer et al¹

	problems, how difficult hav ome, or get along with oth		de it for you to do your work,
☐ Not difficult at all	Somewhat difficult	☐ Very difficult	Extremely difficult
Calculate the patient's a of "not at all," "several c	nxiety severity by assigning lays," "more than half the c	g scores of 0, 1, 2 and days," and "nearly eve	3 to the response categories ery day," respectively.
	seven items ranges from 0 iety, and 15-21 indicates s		indicates mild anxiety, 10-14

Using the threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD (General Anxiety Disorder). It is moderately good at screening three other common anxiety disorders: panic disorder (sensitivity 74%, specificity 81%), social anxiety disorder (sensitivity 72%, specificity 80%), and post-traumatic stress disorder (sensitivity 66%, specificity 81%). When screening for individual or any anxiety disorder, a recommended cut point for further evaluation is a score of 10 or greater.

References: 1. Spitzer RL et al. A brief measure for assessing generalized anxiety disorder. The GAD-7. Arch Intern Med 2006;166:1092-7. 2. Kroenke K et al. Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. Ann Intern Med 2007:146(5):317-25.