Contraindications to and Precautions for Commonly Used Vaccines*

		1	T
Vaccine**	True contraindications	Precautions ++	Not a contraindication
All vaccines	 Anaphylactic reaction to a previous dose of vaccine Anaphylactic reaction to a constituent of a vaccine 	Moderate to severe illness with or without fever	 Mild to moderate local reactions to previous injection of vaccine Mild, acute illness with or without fever Current antimicrobial therapy with the exception of live bacterial vaccines Convalescent phase of an acute illness Prematurity Breastfeeding Recent exposure to infectious disease Personal or family history of allergy
DPT	Anaphylactic reaction to a previous dose of vaccine	Hypotonic- hyporesponsive state within 48 hrs of prior dose of DPT	 Fever > 40.5°C after prior dose of DPT Family history of sudden infant death syndrome Convulsions within 48 hrs. of prior dose of DPT Family history of convulsions Persistent inconsolable crying lasting > 3 hrs within 48 hrs of prior dose Pre-existing neurologic conditions Prior history of pertussis
IPV	Anaphylactic reaction to neomycin		
MMR	 Anaphylactic reaction to previous dose or to neomycin Pregnancy Severe immunodeficiency (See section on Immunization in Immunocompromised Hosts) 	• Recent administration of IG (see Table 7)	 Tuberculosis or positive TB skin test Simultaneous TB skin testing Current antimicrobial therapy Infection with HIV (1994 Pediatric HIV Classification categories E, N1, A1) Egg allergy
Hib			History of Hib disease

-----Table 2 (con't)-----

Contraindications to and Precautions for Commonly Used Vaccines*

Vaccine**	True contraindications	Precautions ++	Not a contraindication
Hepatitis A & B			Pregnancy
Influenza	Anaphylactic reaction to a previous dose, known anaphylactic hypersensitivity to eggs manifested as hives; swelling of the mouth and throat; difficulty in breathing; hypotension and shock		• Pregnancy
Meningococcal vaccines	 Hypersensitivity to any component of the vaccine; history of signs of hypersensitivity after previous administration of the vaccine 		Pregnancy
Pneumococcal vaccines	Hypersensitivity to any component of the vaccine including diphtheria toxoid	Possible history of latex sensitivity	Pregnancy
Varicella	 Immunocompromised people (see Varicella chapter) Pregnancy 	• Recent administration of blood, plasma, IG or VZIG (see Table 7)	 History of contact dermatitis to neomycin Patients with nephritic syndrome or those undergoing hemodialysis and peritoneal dialysis Patients taking low doses of inhaled steroids

^{*} For complete information regarding a particular vaccine, the reader is invited to consult the specific vaccine chapter of the *Guide*.

^{**} DPT = diphtheria, Pertussis and tetanus vaccine. IPV = inactivated poliovirus vaccine. MMR= measles, mumps and rubella vaccine. Hib = Haemophilus influenza type b conjugate vaccine

⁻⁺⁺ The events or conditions listed as precautions are not contraindications, but should be carefully considered in determining the benefits and risks of administering a specific vaccine. If the benefits are believed to outweigh the risks (e.g. during an outbreak or foreign travel) the vaccine should be given.