Adapted with Permission from Appendix G: Recommended Minimum Cleaning and Disinfection Level and Frequency for Non-critical Client/Patient/Resident Care Equipment and Environmental Items

The following chart relates to **non-critical patient care equipment** only, i.e., equipment that comes into contact with intact skin. For semi-critical and critical equipment that require high level disinfection or sterilization, see the Ministry of Health and Long-term Care's *Best Practices for Cleaning, Disinfection and Sterilization in all Health Care Settings*. Refer to Appendix F for appropriate agents that may be used for cleaning and disinfection of non-critical patient care equipment.

This chart also includes **environmental surfaces and items** that do not come into contact with skin. Refer to Section III and Appendix E for guidance regarding cleaning, disinfection of environmental surfaces and items.

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Arrest Cart	See Resuscitation Cart			HCW
Basin (Bath or Wash)	LLD	• After each use	Dry completely before use	HCW

Appendix B

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Bassinette	LLD	WeeklyWhen soiledBetween newborns		HCW
Bed				
Bedrail and extender	LLD	• Daily		ECW
Mattress	LLD	 Clean between patients and when soiled 		ECW HCW – if soiled
Visitor cot	LLD	Change linen and clean between uses		ECW
Bedpan and Urinal				
Single Patient	CL	 Clean after each use if designated to client 	Remove gross soil and fluids before cleaning	HCW
Between patients	LLD	Between clients	Remove gross soil and fluids before cleaning	HCW
Blood Pressure Cuff	LLD	Between clients		LICYA
		When soiled		HCW
Cardiac Monitor	LLD	 Daily and between clients 		HCW

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Cast Cutting Blades Saws	CL or disposable CL	When soiledWhen soiled	Send for sterilization if contact with blood or body fluids	HCW HCW
Chair Includes recliners, and clinic chairs and sofa Chart Cover Binder and/or clipboard	LLD	Daily and when soiled When soiled	 Charts and clipboards should not go into rooms on Additional Precautions 	ECW HCW
Clippers Surgical	LLD	Between clients	Disposable heads are preferred	HCW

Appendix B

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Commode Chairs Single client use	LLD	• When soiled	 Ideally dedicated to each client Clients with VRE or C. difficile, must have dedicated commode For C.difficile, consider cleaning with a sporicidal agent Remove gross soil and fluids before cleaning and disinfection 	ECW
Multiple patient use	LLD	When soiled Between clients	 Remove gross soil and fluids before cleaning and disinfection 	HCW
Defibrillator	See Resuscitation cart			HCW
Diagnostic Imaging Portable-Machine	LLD	When soiled and on leaving Contact Precautions room		HCW
Portable-portable grid/film cassette	LLD	Between clients if not covered	Ideally should be covered (e.g., pillowcase cover)	HCW

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Dopplers Transducers Probes	LLD	After each useAfter each use	 Wipe immediately after use to remove residual ultrasound gel before cleaning Probes that contact mucous membranes or non-intact 	HCW
ECG			skin require high level disinfection	
Machine and Cables	LLD	Between clients		HCW
Examination Table	LLD	Between clients and when soiledDaily cleaning		HCW ECW
Glucometer	LLD	• After each use		HCW
Hydraulic Lift Machine	LLD	• As required		ECW
Sling	Launder	Between clients and when soiled	Dedicated to client if possibleLaunder if visibly soiled	ECW
Intravenous (IV) Pumps, Poles, Warmers	LLD	Between clients When soiled		HCW

Appendix B

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Isolette	LLD	WeeklyWhen soiled		ECW HCW
Laryngoscope Handle Blade Mattress	LLD HLD See Bed	Between clients Between clients		HCW
iviatti ess	See Beu			
Measuring Container (urine) Single client use	CL	• After each use		HCW
Multiple client use	LLD	After each use	One container per client, labelled with name	HCW
Ophthalmoscope	LLD	Between clients		HCW
Orthopedic Equipment Crutches, traction, etc	LLD	Between clients		ECW
Otoscope Handle	LLD	Between clients		HCW
Ear Speculum	Disposable or HLD	• Between clients		HCW

Item	Minimum Cleaning and Disinfection Level:	Minimum Frequency	Remarks	Responsible for Cleaning
	CL=Clean only HLD= Clean + High-level			HCW = Health Care Worker
	Disinfection LLD = Clean+ Low-level Disinfection			ECW = Environmental Cleaning Worker
Oximeter Probes	LLD	Daily and between clients	 If single –use, discard after use Refer to manufacturer's instructions for cleaning 	HCW
Pillow	LLD	Between clients and when soiled	Inspect for cracks and discard as required	HCW
Reflex Hammer	LLD	Between clients		HCW
Restraints	Launder	 Between clients and when soiled 	• Launder	ECW
Resuscitation Cart/Arrest Cart	LLD	Weekly and after use	Avoid taking cart into Contact Precautions room, have a designated clean	HCW
			person to pass supplies as required	HCW
Defibrillator	LLD	After each use	All items taken into Contact Precautions room must be	HCW
Trays	LLD	• After each use	discarded and not returned to the cart, even if unopened	
			 Recommend clear cover if not in use 	

Appendix B

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Scales Adult	LLD	Daily and when soiled		HCW
Diaper Newborn	LLD	After each useAfter each use		HCW HCW
Stretcher	LLD	After each use between clients After single client		HCW ECW
Stethoscope	LLD	After each use	Ideally use own stethoscopeIf shared, disinfect ear pieces	HCW
Suction Machines	LLD	Between clientsWhen soiled		HCW
Table Bedside Over Bed	LLD	DailyWhen soiledBetween clients		ECW HCW
Telementry Equipment Monitor and Cables (e.g., Holter Monitor)	LLD	Between clients When soiled		HCW
Tourniquet	LLD.	Between clients or disposable	Discard when soiled/cracked	HCW
Transfer Boards	LLD	Between clients When soiled		ECW HCW

Appendix B

Item	Minimum Cleaning and Disinfection Level: CL=Clean only HLD= Clean + High-level Disinfection LLD = Clean+ Low-level Disinfection	Minimum Frequency	Remarks	Responsible for Cleaning HCW = Health Care Worker ECW = Environmental Cleaning Worker
Transport Equipment Stretcher Walker Crutches Wheelchair	LLD	Between clientsWhen soiled		ECW HCW
Tub Bath chair	LLD	• After each use		ECW
Ultrasound Transducers Handle and Cable External	LLD	• Between Clients	 Use high-level disinfection for transducer probes if they touch mucous membranes or non-intact skin Recommend cover if not in use 	HCW
Urinal	See bedpan			
Urine Measuring Container	See Measuring Container			
Vacutainer Holder	Single Patient Use			HCW
Walker	See Transport Equipment			
Wall-mounted Oxygen and Suction Fixtures	LLD	Between ClientsWhen soiled		HCW
Water Jug	CL	• Daily	Clean in dishwasher if available	ECW

Wheelchair	See Transport				
	Equipment				