

DOSAGE GUIDE FOR FEVER & PAIN for Canada

DOCTOR RECOMMENDED
#1 Children's
TYLENOL®

RECOMMENDATION FOR:

Infants'
CONCENTRATED
DROPS
80 mg/1 mL



Children's
SUSPENSION
LIQUIDS
160 mg/5 mL



Children's
FASTMELTS®
160 mg per tablet



Children's
Motrin®

Infants'
CONCENTRATED
DROPS
40 mg/1 mL



Children's
SUSPENSION
LIQUIDS
100 mg/5 mL



Age	Weight (Note: For Most Accurate Dosing, Dose By Weight)						
	lbs	kgs					
<input type="checkbox"/> 0-3 mos	Under 12	Under 5.5	Consult a doctor for dosing under 4 months			Consult a doctor for dosing under 4 months	—
<input type="checkbox"/> 4-11 mos	12-17	5.5-7.9	1 mL	Use Infants' TYLENOL® Concentrated Drops		1.25 mL	Use Infants' MOTRIN® Concentrated Drops
<input type="checkbox"/> 12-23 mos	18-23	8-10.9	1.5 mL			1.875 mL	
<input type="checkbox"/> 2-3 yrs	24-35	11-15.9	—	5 mL	1 tablet	—	5 mL
<input type="checkbox"/> 4-5 yrs	36-47	16-21.9	—	7.5 mL	1 ½ tablets	—	7.5 mL
<input type="checkbox"/> 6-8 yrs	48-59	22-26.9	—	10 mL	2 tablets	—	10 mL
<input type="checkbox"/> 9-10 yrs	60-71	27-31.9	—	12.5 mL	2 ½ tablets	—	12.5 mL
<input type="checkbox"/> 11 yrs	72-95	32-43.9	—	15 mL	3 tablets	—	15 mL

Children's TYLENOL® is indicated for the temporary relief of mild to moderate pain in a wide variety of conditions involving musculoskeletal pain, headache pain, earache, the symptomatic reduction of fever due to the common cold, flu and other viral or bacterial infections.

Children's MOTRIN® is indicated for the temporary relief of minor aches and pains in muscles, bones and joints, headache, fever, the aches and fever due to the common cold or flu, immunizations, toothache (dental pain), sore throat and earache. For more information please consult www.tylenol.ca or the package insert.

Infants', Children's & Junior Strength Children's TYLENOL® products dosage guide is based on a 10-15 mg acetaminophen/kg/dose. Single dose may be repeated every 4-6 hours, as needed. It is hazardous to exceed 5 doses of acetaminophen per day. Refer to package insert for dosing instructions. Infants' and Children's MOTRIN® product single doses may be given every 6 to 8 hours as needed, but do not exceed 4 doses per day unless advised by your doctor. Refer to package insert for dosing instructions.

Only Infants' MOTRIN® Concentrated Drops 40 mg/mL and Infants' TYLENOL® Concentrated Drops 80 mg/mL are labelled for children under 2 years of age. Consumer labelling for Children's MOTRIN® Suspension Liquid 100 mg/5 mL, Children's TYLENOL® Suspension Liquid 160 mg/5 mL or for Junior Strength TYLENOL® Fastmelts® Tablets 160 mg does not offer dosing for children under 2 years of age; therefore these doses are provided as a guide for professional recommendations to consumers only. Package label does not provide dosing for use in infants less than 4 months of age for Infants' TYLENOL® Concentrated Drops 80 mg/mL or Infants' MOTRIN® Concentrated Drops 40 mg/mL. Please consult a doctor for dosing under 4 months of age.

© Johnson & Johnson Inc. 2013

WHAT TEMPERATURE INDICATES A FEVER? THAT DEPENDS ON HOW IT WAS TAKEN.

Orally: Higher than 37.5°C (99.5°F)

In the ear: Higher than 38.0°C (100.4°F)

In the rectum: Higher than 38.0°C (100.4°F)

In the armpit: Higher than 37.3°C (99.1°F)

**CALL YOUR DOCTOR IF
YOUR CHILD'S TEMPERATURE
EXCEEDS 38.3°C (101°F)
FOR MORE THAN 24 HOURS.**

For more
helpful advice
visit us on



DOCTOR RECOMMENDED
#1 Infants'
TYLENOL®

FORMULATION DOCTORS
RECOMMEND MOST,
WITH A UNIQUE
MECHANISM DESIGNED
TO IMPROVE
DOSING ACCURACY

NEW SimpleMeasure™ Device



Step 1:
Push air out
of syringe.



Step 2:
Insert syringe
tip into bottle
opening.



Step 3:
Flip bottle
upside-down.



Step 4:
Pull white part
of syringe to
correct dose.



DOSAGE GUIDE FOR COLD, COUGH & FEVER for Canada



PSTY13-21E

NEED MORE
INFORMATION?
1-800-265-7323

RECOMMENDATION FOR:

Infants'
FEVER & SORE
THROAT PAIN



CONCENTRATED DROPS
Per 1 mL
80 mg Acetaminophen

Children's
FEVER & SORE
THROAT PAIN



SUSPENSION LIQUID
Per 5 mL (1 tsp.)
160 mg Acetaminophen

**JUNIOR STRENGTH
Children's**
FASTMELTS®
FEVER & SORE
THROAT PAIN



**CHEWABLE
TABLET**
160 mg Acetaminophen

Children's
COLD & COUGH



SUSPENSION LIQUID
Per 5 mL (1 tsp.):
160 mg Acetaminophen,
15 mg Pseudoephedrine Hydrochloride,
7.5 mg Dextromethorphan Hydrobromide,
1 mg Chlorpheniramine Maleate

Children's
COMPLETE COLD,
COUGH & FEVER



SUSPENSION LIQUID
Per 5 mL (1 tsp.):
160 mg Acetaminophen,
15 mg Pseudoephedrine Hydrochloride,
7.5 mg Dextromethorphan Hydrobromide,
1 mg Chlorpheniramine Maleate

Age	Weight (Note: For Most Accurate Dosing, Dose By Weight)		 CONCENTRATED DROPS Per 1 mL 80 mg Acetaminophen	 SUSPENSION LIQUID Per 5 mL (1 tsp.) 160 mg Acetaminophen	CHEWABLE TABLET 160 mg Acetaminophen	SUSPENSION LIQUID Per 5 mL (1 tsp.): 160 mg Acetaminophen, 15 mg Pseudoephedrine Hydrochloride, 7.5 mg Dextromethorphan Hydrobromide, 1 mg Chlorpheniramine Maleate	SUSPENSION LIQUID Per 5 mL (1 tsp.): 160 mg Acetaminophen, 15 mg Pseudoephedrine Hydrochloride, 7.5 mg Dextromethorphan Hydrobromide, 1 mg Chlorpheniramine Maleate
	lbs	kgs					
<input type="checkbox"/> 0-3 mos	Under 12	Under 5.5	Consult a doctor for dosing under 4 months	—	—	Multi-ingredient cough and cold products: use only in children above 6 years of age	
<input type="checkbox"/> 4-11 mos	12-17	5.5-7.9	1 mL	Use Infants' TYLENOL® Fever & Sore Throat Pain Concentrated Drops			
<input type="checkbox"/> 12-23 mos	18-23	8-10.9	1.5 mL				
<input type="checkbox"/> 2-3 yrs	24-35	11-15.9	—	5 mL	1 tablet	10 mL 2 tablets	10 mL 2 tablets
<input type="checkbox"/> 4-5 yrs	36-47	16-21.9	—	7.5 mL	1 ½ tablets		
<input type="checkbox"/> 6-8 yrs	48-59	22-26.9	—	10 mL	2 tablets		
<input type="checkbox"/> 9-10 yrs	60-71	27-31.9	—	12.5 mL	2 ½ tablets		
<input type="checkbox"/> 11 yrs	72-95	32-43.9	—	15 mL	3 tablets		

FEVER AND DOSAGE DIARY

Use this chart to keep track of your child's fever and medicine dosing. Make sure any other caregivers use it as well.

Date						
Time						
Temperature						
Medicine						
Dose						
Given By						

Always use the measuring device that comes with the medicine.
Always remember to keep package out of reach of children.

CAUTION: If fever persists for more than 3 days, pain for more
than 5 days or new symptoms appear, talk to your doctor.

Infants', Children's & Junior Strength Children's TYLENOL® Fever & Sore Throat Pain single ingredient
acetaminophen do not exceed 5 doses of acetaminophen per day.

Junior Strength and Children's TYLENOL® Cold & Cough and Junior Strength and Children's TYLENOL®
Complete Cold, Cough & Fever Products are not recommended for children under 6 years of age.
Do not exceed 4 doses per day.

HEALTHY ESSENTIALS™
YOUR BEST DAY, EVERY DAY™

Visit the HEALTHY ESSENTIALS™ site for more information on Johnson & Johnson pediatric
products including exclusive coupons, promotions, and helpful lifestyle articles and tools.