

ADULT PAIN AND FEVER RELIEF

For Children's Advil products, see reverse

Advil Liqui-Gels®

200 mg Ibuprofen (per Liqui-Gel)
Analgesic/Antipyretic

For fast and effective relief of:

- Mild to moderate migraine headaches, including associated symptoms of:
 - Nausea – Photosensitivity – Phonosensitivity
- The same indications as Advil Tablets below

Advil



Adults and children over 12:

For migraine headaches, take 1 Liqui-Gel every 4 hours or 2 Liqui-Gels every 6-8 hours at the first sign of symptoms as needed. For all other uses, take 1 Liqui-Gel every 4 hours or 2 Liqui-Gels every 6-8 hours as needed.

Maximum daily dose: 6 Liqui-Gels. Do not use longer than 3 days for a fever or 5 days for pain relief.

Advil Extra Strength Liqui-Gels

400 mg Ibuprofen (per Liqui-Gel)
Analgesic/Antipyretic

For long lasting pain relief of:

- Mild to moderate migraine headaches, including associated symptoms of:
 - Nausea – Photosensitivity – Phonosensitivity
- The same indications as Advil Extra Strength Caplets below

Advil 400



Adults and children over 12:

For migraine headaches, take 1 Extra Strength Liqui-Gel at the first sign of symptoms and every 6-8 hours as needed. For all other indications, take 1 Extra Strength Liqui-Gel every 6-8 hours as needed.

Maximum daily dose: 3 Liqui-Gels. Do not use longer than 3 days for a fever or 5 days for pain relief.

Advil Tablets/Caplets

200 mg Ibuprofen (per tablet/caplet)
Analgesic/Antipyretic

For fast and effective relief of:

- Headache
- Muscle aches
- Joint pain
- Backache
- Menstrual pain
- Toothache
- Fever
- Pain of inflammation

Advil



Adults and children over 12:

Take 1 tablet/caplet every 4 hours or 2 tablets/caplets every 6-8 hours as needed.

Maximum daily dose: 6 tablets/caplets. Do not use longer than 3 days for a fever or 5 days for pain relief.

Advil Extra Strength Caplets

400 mg Ibuprofen (per caplet)
Analgesic/Antipyretic

For long lasting pain relief of:

- Headache
- Muscle aches
- Joint pain
- Backache
- Menstrual pain
- Toothache
- Fever
- Pain of inflammation

Advil 400



Adults and children over 12:

Take 1 caplet every 6-8 hours as needed.

Maximum daily dose: 3 caplets. Do not use longer than 3 days for a fever or 5 days for pain relief.

Advil Arthritis Pain

400 mg Ibuprofen (per Liqui-Gel)
Analgesic

NEW

For fast, long lasting and effective relief of:

- Pain from inflammation associated with arthritis

Advil AP



Adults and children over 12:

Take 1 Liqui-Gel every 6-8 hours as needed.

Maximum daily dose: 3 Liqui-Gels. Do not use longer than 5 days for pain relief.

Advil Muscle & Joint

400 mg Ibuprofen (per caplet)
Analgesic

For long lasting pain relief of:

- Muscle aches
- Joint pain
- Backache
- Menstrual pain
- Headache
- Toothache
- Pain of inflammation

Advil M



Adults and children over 12:

Take 1 caplet every 6-8 hours as needed.

Maximum daily dose: 3 caplets. Do not use longer than 3 days for a fever or 5 days for pain relief.

Advil NIGHTTIME Liqui-Gels

200 mg Ibuprofen • 25 mg Diphenhydramine HCl (per Liqui-Gel)
Analgesic/Sleep Aid

For fast and effective relief of:

- Occasional mild to moderate nighttime pain and accompanying sleeplessness

Advil PM



Age 16 to 65:

Take 1 or 2 Liqui-Gels at night as needed.

Maximum use: 5 days or less.

Advil Day/Night Convenience Pack

Daytime**

200 mg Ibuprofen (per Liqui-Gel)
Analgesic/Antipyretic

For fast and effective relief of:

- Migraines
- Headaches
- Muscle & joint pain
- Backaches

Nighttime***

200 mg Ibuprofen
25 mg Diphenhydramine HCl (per Liqui-Gel)
Analgesic/Sleep Aid

For fast and effective relief of:

- Migraines
- Headaches
- Muscle & joint pain
- Backaches
- Plus a sleep aid

NEW

Advil



Age 16 to 65 (daytime):

Take 1 green Liqui-Gel every 4 hours, or 2 green Liqui-Gels every 6-8 hours at the first sign of symptoms as needed.

Age 16 to 65 (nighttime):

Take 1 or 2 blue Liqui-Gels at night, 4-6 hours after the last daytime ibuprofen or diphenhydramine dose.

Maximum use: Do not exceed a combined total of any 6 green or blue Liqui-Gels in 24 hours. Do not take for more than 5 consecutive days/nights unless directed by a doctor.